arXiv:quant-ph/9704037v1 18 Apr 1997

Number–phase uncertainty relations: verification by homodyning

T. Opatrný[1,2], M. Dakna[1], and D.–G. Welsch[1]
[1] *Friedrich-Schiller-Universität Jena, Theoretisch-Physikalisches Institut*
Max-Wien-Platz 1, 07743 Jena, Germany
[2] *Department of Theoretical Physics, Palacký University*
Svobody 26, 77146 Olomouc, Czech Republic

(April 18, 1997)

It is shown that fundamental uncertainty relations between photon number and canonical phase of a single-mode optical field can be verified by means of balanced homodyne measurement. All the relevant quantities can be sampled directly from the measured phase-dependent quadrature distribution.

42.50.Dv, 42.50.Ar, 03.65.Bz

Although the problem of number–phase uncertainty has widely been studied, there has been no direct experimental verification of fundamental uncertainty relations (URs). What can be the best way of doing that? A powerful and perhaps ultimate method for measuring the quantum statistics of traveling optical fields has been balanced homodyne detection. The quantity that is directly measured is the probability distribution $p(x,\vartheta)$ of the phase-dependent quadrature $\hat{x}(\vartheta)=2^{-1/2}(\hat{a}e^{-i\vartheta}+\hat{a}^{\dagger}e^{i\vartheta})$, where $\hat{a}$ ($\hat{a}^{\dagger}$) is the bosonic annihilation (creation) operator of the (single-mode) signal field and ϑ corresponds to the local-oscillator phase. It has been shown that $p(x,\vartheta)$ for all phases ϑ in a π interval contains all knowable information about the quantum state of the signal field and can be used to reconstruct the Wigner function applying inverse Radon transformation techniques [1].

Since the Wigner function is a full description of the quantum state, it can be used to calculate other important features of the field, such as the photon-number and phase statistics and their associated URs [2]. The inverse Radon transform requires a three-fold integration of the measured data, and the calculation of the density matrix in the Fock basis can then be accomplished with two integrals. One summation eventually yields the photon-number moments, and one sum and one integral must be performed to obtain the (canonical) phase moments. Hence, six- and seven-fold transformations of the recorded data are required for UR verification at least. Of course, large amount of data manipulation accumulates various errors and the physical nature of the uncertainties becomes less transparent.

Recent progress has offered new possibilities of determining the photon-number statistics in a more direct way avoiding the detour via the Wigner function. It has been shown that both the density-matrix elements $\varrho_{nn'}$ in the Fock basis [3] and the moments and correlations $\langle\hat{a}^{\dagger k}\hat{a}^{k'}\rangle$ [4] can directly be sampled from the recorded data according to a two-fold integral transform

$$\mathcal{A}=\int_{2\pi}d\vartheta\int_{-\infty}^{\infty}dx\,K_{\mathcal{A}}(x,\vartheta)p(x,\vartheta). \tag{1}$$

Here, $\mathcal{A}$ is the quantity that is to be determined and $K_{\mathcal{A}}(x,\vartheta)$ is the corresponding integral kernel (sampling function). In particular, $\mathcal{A}$ can be identified with $p_n=\varrho_{nn}$ or $\langle\hat{n}^k\rangle=\langle(\hat{a}^{\dagger}\hat{a})^k\rangle$ in order to determine the photon-number distribution or the photon-number moments.

In contrast to the photon number, the phase has remained a troublesome variable. Recently it has been suggested to measure the canonical phase distribution asymptotically by replacing the local-oscillator with a reference mode prepared in so-called reciprocal binomial states – a method that is state dependent and hardly realizable at present [5]. It has also been suggested to use balanced homodyning for asymptotically sampling phase distributions [6]. Again, the method is state dependent and the exact phase distributions can be obtained only in the sense of a limiting process. Moreover, with increasing (photon-number) excitation of the state under consideration an increasing number of highly oscillating functions must be summed up to obtain the corresponding integral kernels and high-accuracy sampling of the phase distributions from the recorded data becomes very difficult. The problem is similar to that of a calculation of phase quantities from the sampled density-matrix elements in the Fock basis. With increasing excitation of the state a larger number of density-matrix elements must be sampled, so that the accumulated error may eventually dominate the result.

In what follows we show that fundamental number–phase URs that are based on exponential phase measures can be verified directly from the homodyne data according to Eq. (1), without making a detour via other quantities and without making any assumptions and approximations with regard to the state. Let us consider a phase distribution $p(\varphi)$ and its exponential phase moments defined by

$$\Psi_k=\int_{2\pi}d\varphi\,e^{ik\varphi}p(\varphi). \tag{2}$$

In particular for the canonical phase the relation $\Psi_k=\langle\hat{E}^k\rangle$ is valid, where the exponential phase operator $\hat{E}$ is given by $\hat{E}=(\hat{n}+1)^{-1/2}\hat{a}$ (for the canonical phase and the exponential phase operator, see, e.g., [7]). In order to define a mean phase $\bar{\varphi}$ that is independent of the chosen phase window, the first-order exponential phase moment has been introduced in the definition, $\bar{\varphi}=\arg\Psi_1=$

$\arg\langle\hat{E}\rangle$, and it has been used to define phase uncertainty measures, such as

$$\Delta\varphi = \arccos|\Psi_1| = \arccos|\langle\hat{E}\rangle| \tag{3}$$

[8], the Bandilla-Paul dispersion σ_{BP}^2 [9], and the Holevo dispersion σ_{H}^2 [10],

$$\sigma_{\mathrm{BP}} = \sin\Delta\varphi, \quad \sigma_{\mathrm{H}} = \tan\Delta\varphi. \tag{4}$$

It can then be proven that the UR

$$\Delta n \tan\Delta\varphi \geq \tfrac{1}{2} \tag{5}$$

is valid [8], which is equivalent to the Holevo UR [10]

$$(\Delta n)^2 \sigma_{\mathrm{H}}^2 \geq \tfrac{1}{4} \tag{6}$$

$[(\Delta n)^2 = \langle\hat{n}^2\rangle - \langle\hat{n}\rangle^2]$. Although these URs are exact, they are, in a sense, weak. This means that they also allow for such values of Δn and $\Delta\varphi$ for which no state exists (for more specific URs and the corresponding minimizing states, see [11,12]). Further, URs that are based on the Susskind–Glogower trigonometric operators $\hat{C} = \frac{1}{2}(\hat{E} + \hat{E}^\dagger)$ and $\hat{S} = \frac{1}{2i}(\hat{E} - \hat{E}^\dagger)$ have also been studied [7,12,13],

$$\Delta n \Delta C \geq \tfrac{1}{2}|\langle\hat{S}\rangle|, \quad \Delta n \Delta S \geq \tfrac{1}{2}|\langle\hat{C}\rangle|, \tag{7}$$

$$\Delta S \Delta C \geq \tfrac{1}{2}\varrho_{00} \tag{8}$$

$[(\Delta C)^2 = \langle\hat{C}^2\rangle - \langle\hat{C}\rangle^2$, and $(\Delta S)^2$ accordingly]. Note that the squares of the trigonometric operators can be written as

$$\hat{C}^2 = \tfrac{1}{2}\hat{1} + \tfrac{1}{4}\left(\hat{E}^2 + \hat{E}^{\dagger 2}\right) - \tfrac{1}{4}|0\rangle\langle 0|, \tag{9}$$

$$\hat{S}^2 = \tfrac{1}{2}\hat{1} - \tfrac{1}{4}\left(\hat{E}^2 + \hat{E}^{\dagger 2}\right) - \tfrac{1}{4}|0\rangle\langle 0|. \tag{10}$$

From the definitions of $\Delta\varphi$, ΔC and ΔS we see that for measuring them it is sufficient to measure the exponential phase moments $\Psi_1 = \langle\hat{E}\rangle$ and $\Psi_2 = \langle\hat{E}^2\rangle$ and the density-matrix element $\varrho_{00} = \langle|0\rangle\langle 0|\rangle$. The determination of the photon-number uncertainty requires measurement of $\langle\hat{n}\rangle$ and $\langle\hat{n}^2\rangle$.

It is well known that ϱ_{00}, $\langle\hat{n}\rangle$, and $\langle\hat{n}^2\rangle$ can directly be sampled from the data recorded in balanced homodyning applying Eq. (1). The kernel $K_{00}(x,\vartheta)$ for ϱ_{00} can be taken from [3], $K_{00}(x,\vartheta) = \pi^{-1}\,\Phi(1,\frac{1}{2},-x^2)$, where $\Phi(a,b,z)$ is the confluent hypergeometric function. The kernels $K_n(x,\vartheta)$ and $K_{n^2}(x,\vartheta)$ for $\langle\hat{n}\rangle$ and $\langle\hat{n}^2\rangle$, respectively, can simply be obtained from the sampling formula for the normally ordered moments and correlations of bosonic operators [4],

$$\langle\hat{a}^{\dagger n}\hat{a}^m\rangle = \int_{2\pi} d\vartheta\, e^{i(n-m)\vartheta} \int_{-\infty}^{\infty} dx\, \frac{\mathrm{H}_{n+m}(x)}{2\pi\sqrt{2^{n+m}}\binom{n+m}{m}}\, p(x,\vartheta), \tag{11}$$

from which we find that

$$K_n(x,\vartheta) = \tfrac{1}{2\pi}\left(x^2 - \tfrac{1}{2}\right), \tag{12}$$

$$K_{n^2}(x,\vartheta) = \tfrac{1}{2\pi}\left(\tfrac{2}{3}x^4 - x^2\right). \tag{13}$$

Note that for large $|x|$ the leading terms in these expressions determine the kernels for the energy moments in the classical limit.

Let us now turn to the problem of direct sampling of the exponential phase moments of the canonical phase. Provided that the corresponding integral kernels exist, their asymptotic behavior for large $|x|$ can be obtained from considering the classical limit. Since in classical physics the phase probability distribution $p(\varphi)$ can be obtained from the phase-space probability distribution $W(r,\varphi)$ according to $p(\varphi) = \int_0^\infty rdr\, W(r,\varphi)$, the exponential phase moments Ψ_k can be given by

$$\Psi_k = \int_{2\pi}\int_0^\infty rdr\, W(r,\varphi) e^{ik\varphi}. \tag{14}$$

Further, the quadrature probability $p(x,\vartheta)$ is given by the Radon transform

$$p(x,\vartheta) = \int_{2\pi} d\varphi \int_0^\infty rdr\, W(r,\varphi)\, \delta[x - r\cos(\vartheta - \varphi)]\,. \tag{15}$$

Let us now assume that Ψ_k can be related to $p(x,\vartheta)$ according to Eq. (1) [$\mathcal{A} \to \Psi_k$ and $K_{\mathcal{A}}(x,\vartheta) \to K_k(x,\vartheta)$]. Substituting in this equation for $p(x,\vartheta)$ the result of Eq. (15) and comparing with Eq. (14), we observe that $K_k(x,\vartheta)$ can be written as $K_k(x,\vartheta) = e^{ik\vartheta}K_k(x)$, where $K_k(x)$ must satisfy the integral equation

$$\int_{2\pi} d\varphi\, e^{ik\varphi} K_k(r\cos\varphi) = 1 \tag{16}$$

for any $r > 0$. From Eq. (16) we can see that $K_k(x)$ is not uniquely defined. First, any function of parity $(-1)^{k+1}$ can be added to $K_k(x)$ without changing the integral. Second, any polynomial of a degree less than k can also be added to $K_k(x)$. As can be verified, Eq. (16) is solved using the functions

$$K_{2m+1}(x) = \tfrac{1}{4}(-1)^m(2m+1)\,\mathrm{sign}(x) \tag{17}$$

and

$$K_{2m}(x) = \pi^{-1}(-1)^{m+1} m \log|x| \tag{18}$$

($m = 0,1,2,\ldots$). It is worth noting that since Eq. (15) is also valid when $W(r,\varphi)$ is the quantum-mechanical Wigner function, using in homodyne detection the kernels (17) and (18) over the whole x axis would yield the exponential phase moments of the phase quasi-probability distribution defined by the radially integrated Wigner function.

With regard to the canonical phase, the kernels (17) and (18) are of course valid only for large $|x|$. To obtain them for arbitrary $|x|$, we recall that in quantum physics Ψ_k can be given by

$$\Psi_k = \langle \hat{E}^k \rangle = \sum_{n=0}^{\infty} \varrho_{n+k\,n} \tag{19}$$

in place of Eq. (14). Expressing $p(x,\vartheta)$ in terms of the density-matrix elements $\varrho_{nn'}$ as

$$p(x,\vartheta) = \sum_{n,n'=0}^{\infty} e^{i(n'-n)\vartheta} \psi_n(x) \psi_{n'}(x) \varrho_{nn'}, \tag{20}$$

and assuming that Eq. (1) applies to Ψ_k, we again find that $K_k(x,\vartheta) = e^{ik\vartheta} K_k(x)$, but in place of Eq. (16) $K_k(x)$ must now satisfy the integral equation

$$2\pi \int_{-\infty}^{\infty} dx\; K_k(x) \psi_n(x) \psi_{n+k}(x) = 1 \tag{21}$$

for every $n = 0,1,2,\ldots$. Here, $\psi_n(x) = (2^n n! \sqrt{\pi})^{-1/2} \exp(-x^2/2)\,\mathrm{H}_n(x)$ are the energy eigenfunctions of a harmonic oscillator, with $\mathrm{H}_n(x)$ being the Hermite polynomials. From Eq. (21) and the properties of the Hermite polynomials [14] the same arbitrariness in the determination of $K_k(x)$ as in the classical limit is found. To derive an explicit expression, we use the expansion

$$\begin{aligned} \hat{E}^k &= \sum_{n=0}^{\infty} :\frac{\hat{a}^{\dagger\,n} \exp(-\hat{a}^\dagger \hat{a}) \hat{a}^{n+k}}{\sqrt{n!(n+k)!}}: \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{1}{\sqrt{n!(n+k)!}} \frac{(-1)^m}{m!} \hat{a}^{\dagger n+m} \hat{a}^{n+m+k} \end{aligned} \tag{22}$$

(: : introduces normal order) and apply Eq. (11) in order to represent $\langle \hat{E}^k \rangle$ in the form of Eq. (1). Provided that all the correlations $\langle \hat{a}^{\dagger n} \hat{a}^{n+k} \rangle$ exist, we derive

$$K_k(x) = (2\pi)^{-1} \sum_{l=0}^{\infty} C_l^{(k)} \mathrm{H}_{2l+k}(x), \tag{23}$$

where

$$C_l^{(k)} = \frac{(l+k)!}{2^{l+(k/2)}(2l+k)!} \sum_{n=0}^{l} \binom{l}{n} \frac{(-1)^{l-n}}{\sqrt{(n+1)\ldots(n+k)}}\,. \tag{24}$$

In particular, the kernels for sampling the first two moments can be rewritten as, on using standard relations [14],

$$K_1(x) = \pi^{-3/2} x \int_0^{\infty} \frac{dt}{\sqrt{t}\cosh^2 t}\, \Phi\big(2, \tfrac{3}{2}, -x^2 \tanh t\big) \tag{25}$$

(also see [15]), and (apart from an irrelevant constant)

$$K_2(x) = \frac{1}{2\pi} \int_0^{\infty} dt\, \frac{-\mathrm{I}_0(t)}{\cosh^2 t \sinh t}\, \Phi\big(2, \tfrac{1}{2}, -x^2 \tanh t\big) \tag{26}$$

($\mathrm{I}_0(t)$ is the modified Bessel function). Finally, it can be proven that the kernels exist and satisfy the condition (21), i.e., we have found solutions even when the assumption of finite moments fails and Eq. (11) cannot be used.

The kernels $K_1(x)$ and $K_2(x)$ can be evaluated numerically using standard routines. They are plotted in Fig. 1(a) and (b). As expected, they rapidly approach the classical limits given in Eqs. (17) and (18) as $|x|$ increases. Since they differ from the classical limits only in a small region (of a few "vacuum-fluctuation widths") around zero, in practice their evaluation needs applying Eqs. (25) and (26) [or Eq. (24)] only for small values of $|x|$, whereas for greater values the expressions given in Eqs. (17) and (18) can be used. Note that for small values of $|x|$ power series expansion of the confluent hypergeometric function in Eqs. (25) and (26) can be used.

To verify URs connected with $\Delta\varphi$ [e.g., the relations (5) and (6)], the first moment of $\hat{E}$ must be measured, which can be accomplished with the kernel $K_1(x,\vartheta)$. With regard to URs of the type given in Eqs. (7) and (8), one also needs the second moment of $\hat{E}$ and the vacuum probability ϱ_{00}. From the above, the kernels for sampling $\langle \hat{C}^2 \rangle$ and $\langle \hat{S}^2 \rangle$ read as

$$K_\pm(x,\vartheta) = \tfrac{1}{4\pi}\big[1 - \Phi(1, \tfrac{1}{2}, -x^2)\big] \pm \tfrac{1}{2} \cos(2\vartheta) K_2(x) \tag{27}$$

[see Fig. 1(c)], where $K_+(x,\vartheta) = K_{C^2}(x,\vartheta)$ and $K_-(x,\vartheta) = K_{S^2}(x,\vartheta)$.

To conclude, we have presented a method for verification of number–phase URs. It is based on the possibility of direct sampling of exponential phase moments of the canonical phase of a single-mode quantum state, $\Psi_k = \langle \hat{E}^k \rangle$, from the data recorded in balanced homodyning. We have shown that the corresponding kernels $K_k(x,\vartheta)$ are state independent and well behaved. With increasing $|x|$ they rapidly approach the classical limits. Since the method does not only apply to the determination of low-order moments, it may also be used for reconstructing the canonical phase distribution $p(\varphi)$ as a whole. However, a direct (state independent) sampling of $p(\varphi)$ according to Eq. (1) seems to be impossible. If there would exist a corresponding kernel, its Fourier components with respect to ϑ would be equal to the kernels for determining Ψ_k. Since for chosen x the absolute values of these kernels increase with k [cf. Eqs. (17) and (18)], they cannot be treated as Fourier coefficients of a well-behaved function of ϑ. Of course, this does not exclude an indirect reconstruction of $p(\varphi)$. Measuring a limited number of Ψ_k, one can use, e.g., the maximum entropy principle [16] to obtain a $p(\varphi)$ which best fits the measured values without introducing any arbitrary bias. This also offers the possibility of verification of URs that are not based on exponential phase moments [17,18].

This work was supported by the Deutsche Forschungsgemeinschaft.

---

[1] K. Vogel and H. Risken, Phys. Rev. A **40**, 2847 (1989); D.T. Smithey, M. Beck, M.G. Raymer, and A. Faridani, Phys. Rev. Lett. **70**, 1244 (1993); D.T. Smithey, M. Beck, J. Cooper, M.G. Raymer, and A. Faridani, Phys. Scr. T**48**, 35 (1993).

[2] M. Beck, D.T. Smithey, and M.G. Raymer, Phys. Rev. A **48**, R890 (1993); D.T. Smithey, M. Beck, J. Cooper, and M.G. Raymer, Phys. Rev. A **48**, 3159 (1993).

[3] G.M. D'Ariano, C. Macchiavello, and M.G.A. Paris, Phys. Rev. A **50**, 4298, (1994); U. Leonhardt, H. Paul, and G.M. D'Ariano, Phys. Rev. A**52**, 4899 (1995); M. Munroe, D. Boggavarapu, M.E. Anderson, and M.G. Raymer, Phys. Rev. A **52**, R924 (1995); S. Schiller, G. Breitenbach, S.F. Pereira, T. Müller, and J. Mlynek, Phys. Rev. Lett. **77**, 2933 (1996); U. Leonhardt, M. Munroe, T. Kiss, Th. Richter and M.G. Raymer, Opt. Commun. **127**, 144 (1996); Th. Richter, Phys. Lett. **211**A, 327 (1996).

[4] Th. Richter, Phys. Rev. A **53**, 1197 (1996).

[5] S.M. Barnett and D.T. Pegg, Phys. Rev. Lett **76**, 4148 (1996).

[6] M. Dakna, L. Knöll, and D.-G. Welsch, Phys. Rev. A **55**, 2360 (1997); Quantum Semiclass. Opt., to be published.

[7] R. London, Z. Physik **37**, 915 (1926), **40**, 193 (1927); L. Susskind and J. Glogower, Physics **1**, 49 (1964); P. Carruthers and M.M. Nieto, Rev. Mod. Phys. **40**, 411 (1968).

[8] T. Opatrný, J. Phys. A **27**, 7201 (1994).

[9] A. Bandilla and H. Paul, Ann. Phys. (Lpz) **23**, 323 (1969).

[10] A.S. Holevo, Springer Lecture Notes in Math. **1055**, 153 (1984).

[11] A. Bandilla, H. Paul, and H.-H. Ritze, Quantum Opt. **3**, 267 (1991); A. Bandilla, Optics Commun. **94**, 273 (1992); A. Bandilla and H.-H. Ritze, Phys. Rev. A **49**, 4912 (1994); T. Opatrný, J. Phys. A **28**, 6961 (1995).

[12] A. Lukš and V. Peřinová, Czech. J. Phys. **41**, 1205, (1991); Phys. Rev. A **45**, 6701 (1992); M. Freyberger and W. Schleich, Phys. Rev. A **49**, 5056 (1994).

[13] R. Jackiw, J. Math. Phys. **9**, 339 (1968); M. Kitagawa and Y. Yamamoto, Phys. Rev. A **34**, 3974 (1986); M. Kitagawa, N. Imoto, and Y. Yamamoto, Phys. Rev. A **35**, 5270 (1987). J.H. Shapiro and S.R. Shepard, Phys. Rev. A **43**, 3795 (1991).

[14] A.P. Prudnikov, Yu.A. Brychkov and O.J. Marichev, *Integral and Series*, Vol 2 (Gordon and Breach, New York, 1986).

[15] G.M. D'Ariano, *Homodyning as universal detection,* in: Quantum Communication, Computing and Measurement. Eds. O. Hirota, A.S. Holevo, and C.M. Caves (Plenum, 1997); quant-ph/9701011.

[16] E.T. Jaynes, Phys. Rev. **106**, 620 (1957).

[17] J.A. Vaccaro and D.T. Pegg, J. Mod. Opt. **37**, 17 (1990); S.G. Summy and D.T. Pegg, Optics Commun. **77**, 75 (1990); A.A. Kulaga and F.Ya. Khalili, Zh. Exp. i Teor. Fiz. **104**, 3358 (1993); A. Lindner, D. Reiss, G. Wassiliadis, and H. Freese, Phys. Lett. A **218**, 1 (1996).

[18] I. Białynicki-Birula and Y. Mycielski, Commun. Math. Phys. **44**, 129 (1975); I. Białynicki-Birula, M. Freyberger, and W. Schleich, Phys. Scr. **T 48**, 113 (1993).



FIG. 1. The kernel for sampling the first (a) and second (b) exponential-phase moment is shown (full line) and compared with the classical result (dashed line). The kernel for sampling $\langle\hat{C}^2\rangle$ (c) is shown for $\vartheta = 0$ (1), $\vartheta = \pi/4$ (2), and $\vartheta = \pi/2$ (3).